



This is a repository copy of *Spatial organisation of drumlins*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/118197/>

Version: Accepted Version

Article:

Clark, C.D. orcid.org/0000-0002-1021-6679, Ely, J.C., Spagnolo, M. et al. (3 more authors) (2017) Spatial organisation of drumlins. *Earth Surface Processes and Landforms*. ISSN 0197-9337

<https://doi.org/10.1002/esp.4192>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Spatial organisation of drumlins

[Accepted version - 19 June 2017 - for *Earth Surface Processes and Landforms*]

Chris D. Clark^{1*}, Jeremy C. Ely¹, Matteo Spagnolo², Ute Hahn³, Anna L.C. Hughes⁴ and Chris R. Stokes⁵

¹Department of Geography, University of Sheffield, Sheffield, S10 2TN, UK,
²Department of Geography and Environment, School of Geosciences, University of Aberdeen, Aberdeen, UK, ³Department of Mathematics, Aarhus University, Denmark, ⁴Department of Earth Sciences, University of Bergen and Bjerknes Centre for Climate Research, Bergen, Norway ⁵ Department of Geography, University of Durham, Durham, UK.

*e-mail: c.clark@sheffield.ac.uk

Abstract:

Ice-sheets flowing over soft sediments produce undulations in the bed, typically of metres in relief, of which drumlins are the most abundant and widely investigated. Consensus regarding their mechanism of formation has yet to be achieved. In this paper we examine the spatial organisation of drumlins in order to provide an improved description of the phenomenon and to guide hypotheses of their formation. We review the literature highlighting contradictory findings regarding drumlin spatial organisation and then use this to motivate our study

24 *based on a large sample (42,488) of drumlins from Canada, Britain and Norway.*
25 *Are there typical arrangements in drumlin positioning and are they organised in a*
26 *regular spatial manner (patterned) or are they distributed randomly? We*
27 *recognise that drumlin fields are inherently patchy and therefore apply*
28 *inhomogeneous spatial statistics in order to study their distribution. This shows*
29 *that whilst drumlins are occasionally randomly placed, their main state is non-*
30 *random. They exhibit a strong and statistically significant signal of regularity*
31 *across lengths scales of 100 – 1200 m. We conclude that patterning is a near*
32 *ubiquitous property of drumlins. This finding of regularity demonstrates spatial*
33 *self-organisation in the bedforming process with drumlins as an emergent*
34 *manifestation of subglacial sediment mobility. Kilometre-scale interactions*
35 *between drumlins must occur as they evolve, or interactions may arise as a*
36 *consequence of growth or migration. Hypotheses or models are required that*
37 *can explain the regular spacing of drumlins. We highlight three suggestions for*
38 *such self-organisation: instability in the coupling of ice flow - sediment flux - bed*
39 *shape; local feedback between sediment mobility and relief; and coarsening by*
40 *growth or migration.*

41
42
43 *Key words: Drumlins, patterning, regularity, subglacial bedforms, self-*
44 *organisation.*

Introduction

Somewhat counter-intuitively, ice-sheets abhor flat beds when flowing over soft sedimentary substrates. They produce undulated surfaces, metres to a few tens of metres in relief and with length-scales of hundreds of metres to kilometres and that have been categorised into a number of easily recognised states with well-known geomorphological labels; those oriented transverse to flow (ribbed or Rogen moraine; Dunlop and Clark, 2006, to mega-scale transverse ribs; Greenwood and Kleman, 2010) or parallel to flow and varying in length, from drumlins (Clark *et al.*, 2009; Eyles *et al.* 2016) to mega-scale glacial lineations (Clark, 1993; Spagnolo *et al.*, 2014). They can be thought of as a family of landforms called subglacial bedforms (Aario, 1977; Rose, 1987; Ely *et al.*, 2016) with an implied genetically related formation process, or each type (drumlin, ribbed moraine etc.) or sub-type (e.g. ribbed moraine; Rogen moraine; Blattnick moraine; minor ribbed moraine etc. Hattestrand, 1997) can be regarded as fundamentally different landforms that were formed by different processes.

These subglacial landforms are estimated to cover around half the area previously occupied by palaeo-ice sheets (Clark *et al.*, 2009), with drumlins by far the most abundant type. Drumlins have recently been discovered beneath the Antarctic Ice Sheet (King *et al.* 2009) and emerging from an Icelandic ice cap (Johnson *et al.* 2010; Jónsson *et al.* 2014).

Consensus regarding the formation of drumlins has yet to emerge in spite of two centuries of investigation (Hall, 1815). Rather, there are many hypotheses, ideas

70 and conceptual models, seeking to explain their internal sedimentary structure
71 and the shape, size and pattern of the landforms (overviews by Menzies, 1979;
72 Patterson and Hooke, 1995; Shaw, 2002; Clark, 2010; Stokes et al. 2011; Eyles
73 *et al.* 2016). Progress has matured with some model-based development of the
74 physics of ice-sediment-water interactions at the base of ice sheets (Hindmarsh,
75 1998; Fowler, 2000; Iverson 2000; Hooke and Medford 2013), but these have yet
76 to be developed to yield many predictions for testing, although see Dunlop *et al.*
77 (2008) , Fowler *et al.* (2013) and McCracken et al. (2016) for such model-data
78 comparisons. As with many seemingly intractable scientific problems,
79 researchers from various disciplines (geomorphologists, glaciologists,
80 sedimentologists, and theoreticians) have approached the task from different
81 angles, leading to useful observations, theory and insights and some
82 disagreement regarding how these connect. The so-called ‘drumlin problem’
83 (Menzies and Rose, 1987) remains difficult, and we do not solve it here. Our
84 approach in this paper is to make progress by investigating the spatial
85 organisation of drumlin positioning within their flow-sets; are there typical
86 arrangements and are drumlins organised in a regular spatial manner or are they
87 distributed randomly? The former is indicative of an overall organisation within
88 drumlin flow-sets, suggesting they are a patterned phenomenon, of which nature
89 provides numerous examples (Ball, 1999). Conversely, the latter is consistent
90 with a ‘site-specific’ view whereby each drumlin forms in isolation from its
91 neighbours. Data and findings answering these questions should guide the
92 development of theory or could be used to test numerical process-models that

are both starting to make quantitative predictions and to yield three dimensional model simulation movies of bedform organisation, evolution and migration (Chapwanya et al 2011; Barchyn et al. 2016). Do we need to explain the development of the individual bump (the drumlin) or the wavelength (spacing) and relief of the undulating surface; the drumlin field? Does the overall bumpiness arise in a single phase of development (triggered simultaneously, like measles) or arise from incremental additions of new bumps that 'know of' the existing ones? Rippled surfaces created on a beach in one tide would be an example of the former and Barchan dunes growing and migrating as individuals across a desert are the latter. Unlike these non-glacial examples, we have yet to observe a flat surface beneath an ice sheet grow into drumlins, thus allowing many questions to remain, and highlighting why investigating spatial properties of the final forms might be useful. We review the literature to highlight observations and analyses regarding drumlin spatial organisation and then use these to motivate our study based on a large sample of drumlins.

Existing observations and analyses on spatial organisation

The literature on drumlins abounds with comments on the relationship of a drumlin to its neighbours (see reviews by Menzies, 1979; Patterson and Hooke, 1995; Clark 2010; Eyles et al 2016). An *en echelon* arrangement (Figure 1A) has often been mentioned (e.g. Armstrong and Tipper, 1948; Clapperton, 1989) although a survey of such findings (Patterson and Hooke, 1995) notes that it is not common and no field of drumlins has been found where such positioning

dominates. Much more common is the observation of transverse banding of drumlins (Figure 1B) such that in the downstream direction there are alternating bands of high and low densities. Hill, (1973) for example quantified such variations for ca. 3900 drumlins in Ireland. Elsewhere, the size and shape of drumlins within each band have been reported to be similar (Shaw and Kvill, 1984).

The spacing between neighbouring drumlins is a useful measure that quantifies an aspect of their overall spatial organisation; a preference for a specific spacing implying regularity (Figure 1D). Typically, drumlins are usually found to be widely-spaced with large gaps between them, but in places they have also been reported as tightly packed (i.e. centre-to-centre spacing controlled by drumlin size; Rose and Letzer, 1977) as depicted in Figure 1C. Reed *et al.* (1962) reported 204 measurements of drumlin-to-drumlin spacing in three drumlin fields in the USA, finding their spacing to be rather varied (and multimodal) but with some tendency towards periodicity (i.e. a single preferred spacing). With a larger sample size of drumlins (733) in Ireland, Vernon (1966) found no preferred spacing and that they varied throughout the fields investigated. Drumlin spacing in Poland (Baranowski, 1969) and Canada (Clark and Wilson, 1994) was again found to be varied, but in Ireland and with a larger sample size of 3900, Hill (1973) reported a preferred spacing of around 300 m. Consistent with the above papers, and according to the review by Menzies (1979), for drumlins sampled in the USA, Canada, Ireland and Poland the frequency distributions of spacing were

sometimes found unimodal and normally-distributed, or multimodal and with wide variations in average spacing leading to the conclusion of no overall preferred spacing for drumlins. It is possible however, that if any actual preferred spacing of drumlins exists that some combination of varied mapping approaches, or non-representative sample sizes might be masking the answer. Or perhaps, genuine differences in preferred spacing exist from place to place whose reasons (substrate geology or glaciological controls?) could be usefully sought.

The arrangement of drumlins within flow-sets may provide clues for theories of drumlin formation; inferences of 'process from pattern'. Are they positioned randomly or as regular arrays or as a series of clusters (Figure 1D)? Random positioning appears to have been the most commonly presumed arrangement, in part promoted by the nearest neighbour statistical analyses of Smalley and Unwin (1968), and which may have steered theory development towards drumlins developing from (randomly distributed) sediment inhomogeneities (e.g. patches of stiff till or gravel) or from bedrock bumps. Francek (1991) using a much larger sample (>4000) than the 1968 investigation also reported drumlins to be distributed mostly randomly when analysed by statistical analyses. Contrary to these statistically-based findings, many have remarked on the apparent regularity expressed by drumlins (e.g. Werth, 1909 cited in Baranowski, 1969; Baranowski, 1977; Carl, 1978; Smalley and Warburton, 1994; Fowler, 2000; Clark, 2010) thereby arguing for some form of regularising process inherent in drumlin formation. Examples of drumlins that appear to exist with a repetitive,

systematic arrangement are not difficult to find (e.g. Figure 2) and from such observations many might simply infer that drumlins are positioned regularly. However, as Table 1 highlights published investigations involving statistical analyses of drumlin distribution shows that there is no consensus regarding whether drumlins are placed regularly, randomly, or clustered. Different areas and statistical techniques have produced all three results.

Methods

Data set

In order to address the uncertainties in drumlin spatial organisation highlighted above we used a GIS database comprising 42,488 drumlins spread across 72 different drumlin fields, three palaeo-ice sheets and countries, and across a wide diversity of geological substrates. Most of these (36,222) were mapped across Britain from a high resolution (5 m) digital elevation model (Hughes *et al.*, 2010). Additional mapping was conducted from Landsat ETM+ imagery (15 m resolution) in Alta, Norway (1,483) and Ungava Bay, Canada (5,903). An advantage of this database over those used in previous studies is that it has a larger sample size, covers three palaeo ice sheets, and uses drumlins from highly geologically diverse locations such that drumlins in the sample cover a range of geological substrates and with widely varying drift thicknesses, although we do not analyse these here. See Greenwood and Clark (2010) for such an analysis in Ireland.

185

186 The dataset (Figure 3) was divided into flow-sets (i.e. fields interpreted as
187 forming under single phases of ice flow) based upon their morphology, parallel
188 conformity, spacing and orientation (Clark, 1999).

189

190 *Transverse banding*

191 We used our database to address Hill's (1973) query as to whether the
192 transverse banding he reported for some Irish drumlins is a ubiquitous
193 characteristic. A simple procedure manually linking (by drawing lines in the GIS)
194 laterally adjacent drumlins was adopted. The rules being that an adjacent drumlin
195 belongs to a transverse band if it is positioned laterally, is of similar size and
196 orientation and is not further away than around four times the local drumlin width.
197 This latter rule avoids linking drumlins that are too distant. A quantitative rule-
198 based GIS algorithm could no doubt be devised to accomplish this task with less
199 potential for bias, but we judge our manual method appropriate to a
200 reconnaissance level exploration of the extent to which transverse banding
201 exists. Of arguably greater concern than any such bias is that randomly
202 positioned objects will exhibit alignments by chance and we seek to know if there
203 is any spatial organisation above this level. Figure 4 illustrates this issue by
204 comparing a real drumlin field to a simulated one whereby ellipses of equivalent
205 size and orientation have been randomly positioned across the same domain.

206

207 *Nearest neighbour distances*

To assess any tendency for drumlins to exist at a preferred spacing (see Figure 1D) nearest neighbour distances were calculated as Euclidean distance (within software ArcGIS) between the centroids defined for each drumlin. Centroids were calculated automatically in ArcGIS as the geometric centre of the drumlins, i.e. the arithmetic mean position of all the points in the drumlin's polygon shape. Across-flow spacing of drumlins was also calculated using the mean ice-flow direction (from 10 neighbouring drumlins) and measuring the shortest orthogonal spacing between the centroids of adjacent drumlins. Measurements were used to derive frequency histograms of drumlin spacing.

Spatial statistical methods for assessing regularity.

In order to investigate the nature of spatial organisation in the distribution of drumlins (random, clustered, regular) statistical methods are required, but with spatial phenomena this is not trivial (Kulldorff, 2006). Tests for randomness, such as the Clark-Evans statistic used in the first published investigation of drumlin distribution (Smalley and Unwin, 1968) and the more advanced Ripley's K and L functions (Ripley, 1988), compare measured distances between drumlins to expectations of complete spatial randomness (CSR) producing a metric indicative of dispersion (regularity), randomness or clustering. The K-function ($K(r)$) is:

$$K(r) = \frac{\lambda^{-1} \sum_{i \neq j} (e_{ij} < \pi r^2)}{n}$$

whereby λ is the mean density of points, e_{ij} is the Euclidean distance between the i^{th} and j^{th} points in the dataset, r is the radius of the search area and n is the number of points within the dataset. With CSR, the expected value is:

$$(1) \quad K(r) = \pi r^2$$

For data analysis the related L-function is more appropriate to use as it is more stable and is defined as:

$$(2) \quad L(r) = \sqrt{K(r)/\pi}.$$

These functions have been applied in numerous fields such as ecology and cell biology (Dixon, 2012). Unfortunately, these homogenous tests are sensitive to the area being tested, and do not consider changes in point density across a domain. This is an acute problem for drumlins because we have no *a priori* knowledge of the precise spatial domain. Should this be defined by a convenient box or by the limits of the furthest drumlins? For the latter there is rarely a clear edge, just a decrease in density. It is noteworthy that this issue of sensitivity to the chosen domain led to the later retraction (Unwin, 1997) of the original and widely assimilated conclusion of drumlins having a random distribution (Smalley and Unwin, 1968).

A further problem for any analyses is that areas exist within drumlin fields where drumlins are absent (e.g. Figure 2b). Should these gaps be regarded as 'no observations' and thus be included or as 'no data' because drumlinisation could

not occur here and so should not be part of the domain? Such absences often exist due to a lack of sediment cover, although preservation and visibility issues caused by lakes, bogs and urban areas also occur.

We undertook extensive experimentation using Ripley's K ($K(r)$) and L ($L(r)$) functions for analysing spatial point patterns. We used these functions (in R statistical software) to compare measured distances between drumlins to expectations of complete spatial randomness (CSR), producing a metric indicative of dispersion (regularity), randomness or clustering. To further this approach it is common to use simulation envelopes to compare the observed result to Monte-Carlo simulations of CSR derived from a Poisson (or other) process (Baddeley *et al.*, 2014). Defining the bounding box is an important consideration, as it forms the spatial domain over which simulations of CSR are implemented and thus influences the null hypothesis to be tested. We used a Ripley-Rasson bounding box, which is a convex hull enlarged by a factor dependent upon the number of points within the study area (Ripley and Rasson, 1977). Our analyses across the drumlin database led to confusing results, reported in Ely (2015). By varying the domain size over which the function is performed, we found it possible to yield different answers as to the spatial organisation of drumlins, with clustering as the most common conclusion (and see Maclachlan and Eyles 2013). Typically, with the domain encompassing the whole flow-set, the results tend towards clustering, but if a reduced domain is positioned over a prominent patch of higher density drumlins then regularity is

often revealed such as in Figure 5, but the sample size for such an assessment is low.

It is this seemingly contradictory finding that led the realisation that drumlin fields are actually more appropriately characterised as patches of high density within a background of lower density and with many large gaps (Figure 6A). Such inhomogeneity turns out to be critical with regard to how the distributions are analysed. By way of explanation, if Ripley's functions are performed on points on a regularly spaced grid (with obvious regularity) but with areas rubbed out (Figure 6B) the dominant result will be clustered, even though we can clearly see that there is strong regularity within the clusters. We suspect that the widely varied results reported in the literature (see Table 1) is a consequence of drumlin fields typically being patchy and the methods used being overly sensitive to this patchiness. This issue of inhomogeneity has been widely recognised in ecology, where variables such as soil availability and nutrition introduce inherent clustering of tree positioning (Perry et al., 2006) that is independent of any biophysical spacing controls between individual trees. If we are not to be misled, a technique is therefore required which accounts for the inherent patchiness within a drumlin flow-set; a method of spatial analysis for inhomogeneous point patterns rather than homogenous patterns is required. This is important because arbitrarily defining clusters can be problematic (Couteron et al., 2003) and if one experiments using the Ripley method within individual clusters (e.g. Fig. 5) then the sample sizes typically become too low to gain meaningful results.

Furthermore, Ripley L is a cumulative function, accounting for all previous measurements in subsequent iterations of the search radius. Thus, inferences at specific scales are difficult to make. Therefore, a different method is required for testing whether spatial positioning of drumlins is regular, random or clustered because they have inhomogeneous spatial distributions.

Inhomogeneous Pairwise Correlation Function (g-inhom)

Ripley's K-function and the L-function are cumulative in the sense that K(r) and L(r) count all points (drumlins) with distance at most r to a randomly selected point of the pattern. If one wants to infer at specific distances r, one should use the pair correlation g(r) function instead (e.g. Perry et al., 2006). It is derived from Ripley's K-function in the following manner:

$$g(r) = \frac{1}{2\pi r} \frac{\delta K(r)}{\delta r}$$

The pair correlation function tells us how likely it is to encounter another point of the pattern at a distance r from a 'typical' point, as compared to the corresponding likelihood under complete spatial randomness. Thus, g(r) = 1 means that, seen from a typical point of the pattern, the world at distance r looks completely random (i.e. like a Poisson point process), implying no interactions between points. If, however, g(r) < 1, then points at distance r are less likely to occur than they would under CSR; this is termed 'repulsion' between points at distance r, or that points have been inhibited from occurring at this scale.

Conversely, $g(r) > 1$ means ‘attraction’ between points at distance r ; they are more likely to occur than under CSR. Regular point patterns are characterized by repulsion at small r , followed by a peak of attraction at the typical neighbour distance, while clustered point patterns show attraction already for small values of r (Illian *et al.*, 2008, p. 240).

The original pair correlation function is as sensitive to patchiness and choice of domain boundaries as is the K-function. However we will see that this problem can be overcome by using the inhomogeneous pair correlation function, $g\text{-inhom}$ (Baddeley *et al.*, 2000; Figure 6d). The $g\text{-inhom}$ function is thus appropriate for deciphering the spatial distribution of drumlins. In order to account for inhomogeneities in a dataset, the function takes into account the intensity (density) of the observed points. This is advantageous over arbitrarily defining smaller regions of perceived homogeneity and then performing a homogenous test (Couteron *et al.*, 2003). Instead, $g\text{-inhom}$ weights the point counts according to the local intensity of points within the observed pattern. It is equivalent to comparing the observed pattern with inhomogeneous CSR of same (inhomogeneous) intensity.

In order to assess the spatial arrangement of drumlins over a long range, we computed $g\text{-inhom}(r)$ for a range of r -values. We then simulated CSR through Monte Carlo simulations of a Poisson process producing the same number of observations and point density, to derive a significance envelope (Baddeley et

al., 2014). Deviation from this envelope indicates that the observed patterns are different from CSR. Recently, a technique for assigning significance to this deviation has been developed (Myllymäki et al., 2016), and is implemented here.

The above approach was implemented in the open-source statistical software R. We employed a Ripley-Rasson bounding box (Ripley and Rasson, 1977) and the same edge corrections as used for the Ripley L tests performed above. The libraries 'spatstat' (Baddeley et al., 2015) and 'spptest' (Myllymäki et al., 2016) were used to perform the tests and generate Monte-Carlo significance envelopes. A typical example output from the procedure and labelling is shown in Figure 7.

To account for the clustering in inhomogeneous point patterns an estimation of the point intensity (density) is required. In spatstat, this is done by a kernel density estimator yielding a pixel image. The intensity value in a given pixel is obtained as the weighted number of points within a given search radius. This distance is also called the "kernel band width". As a parameter of the estimation procedure, the search radius strongly influences the resulting intensity estimate (O'Sullivan and Unwin, 2010, p.70), and thus the estimate of g-inhom. Therefore, sensitivity analysis was conducted upon this parameter in order to determine an appropriate value for analysis. Flow-set 29 in southern Scotland (Hughes, *et al.* 2010; 2014) was chosen for sensitivity analysis due to the high number of drumlins ($n = 1,473$). Figure 8 reports the sensitivity of g-inhom(r) to

perturbations of the default value (as a %) of the search radius. A selection of results (80%, 40%, 30%, 5 %, Figure 8) are presented but were explored at 10 % increments between 100 and 10 % and then also at 5%. For the default value (100%) the intensity of points across the flow-set is smooth increasing away from the edges of the flow-set due to edge corrections and appears similar to the 80% case shown in Figure 8. Small scale dispersion is apparent, followed by a strong peak in attraction and final decay to CSR in a manner similar to Figure 7. This is the typical signal across most values in the sensitivity analysis (100 – 30%). With further reduction in the search radius (lower %) clusters within the data become apparent (column two of Fig 8). Reduction of the search radii below 30% of the default value increases the size of the CSR envelope at smaller values (< 500 m). By 5% of the default value the peak no longer occurs above CSR, as the intensity estimate forces simulations of CSR to occur close to the drumlin pattern, almost mirroring it. However, at 5% the intensity map has identified almost as many clusters as there are drumlins such that often only a single drumlin or pair of drumlins is within a single intensity cluster and this is clearly below the length-scale at which we are investigating spatial organisation. For all other flow-sets, a similar pattern of an initial smoothing across the flow-set with an increasing clustering at 30% is apparent. Thus, 30% of the default bandwidth parameter was chosen for all flow-sets.

Results

Transverse banding

Banding is found to be widespread, with 48% of the drumlins analysed for banding (the British examples, $n=36,222$) found to belong to a transverse band (Figure 9). This extends the findings of Hill (1973) showing that such banding is a common feature of the drumlins we have examined and shows that there is a strong element of spatial organisation in drumlin positioning. The mean downstream spacing of bands ($n=1,885$) is 2026 m and the frequency distribution approximates that of a lognormal distribution with a mode of around 800 m. Although we do not demonstrate it here, our visual examination of drumlins elsewhere in the world (using Google Earth) suggests that transverse banding is commonplace. We note that our definition of what constitutes a band is subjective as was our visual method of defining them, so the results quantifying the spacing should be taken as indicative only, pending further investigation. Below we report quantitative and more robust investigations on spatial organisation.

Nearest neighbour spacing

The nearest neighbour spacing of drumlins (Figure 10; British drumlins) reveals a strongly preferred concentration - a uni-modal peak - around 200 to 450 m. This provides a strong hint that regularity exists, but is not proof because relative spatial positions are not analysed and the preferred spacing could be a consequence of drumlin dimensions. A second analysis, designed to assess for the effect of drumlin size was performed; is the preferred spacing simply a consequence of drumlins being packed into a limited area (i.e. Fig 1c)? The

analogy here is how ball-size in a container full of balls controls their spacing (i.e. spatial inhibition). Nearest neighbour analysis conducted orthogonal to ice flow direction revealed drumlin spacing (centroid to centroid; mean of 386 m) to exceed drumlin width (mean of 226 m). This rules out drumlin dimensions being the primary control on spacing. Visual inspection of the arrangement of typical drumlin fields (e.g. Figure 2) leads to the same conclusion in that they are usually found to be widely-spaced rather than packed together and touching.

Spatial statistical methods for assessing degree of regularity

When applied to a typical flow-set the plot of $g\text{-inhom}(r)$ exhibits a clear signal of attraction, well above that which occurs due to random fluctuations and with a prominent peak at 330 m (Figure 11). This demonstrates that there is a strong and statistically significant attraction of points within this nearest neighbour distance, between the length scales of 160 - 1250 m. This combination of short range repulsion followed by a strong peak in attraction is typical of regular patterns. Moreover, of the 42,488 drumlins analysed, 74% of the flow-sets, including those in Canada and Norway, displayed regularity as summarised in Fig 12, representing 93% of the drumlins. By accounting for inherent patchiness and gaps in drumlin fields, this is the first robust demonstration that regularity exists and that it is the predominant signal of drumlin spatial organisation. Furthermore, those drumlins that were found to be not regularly distributed tended to occur in flow-sets that contained small numbers of drumlins ($n < 300$). The attraction peak of $g\text{-inhom}(r)$ occurs at a mean value of ~ 600 m, with the

curve often continuing to 2000 m, indicating that drumlins prefer to be positioned a non-random distance apart across this length scale. Such regularity demonstrates drumlins to be a patterned phenomenon.

Discussion

The finding of widespread banding and a uni-modal peak in the nearest neighbour analyses suggest that drumlins are not randomly placed, but these simple approaches are not a robust demonstration of this. However, by correctly accounting for the patchiness and gaps in drumlin fields by using an appropriate statistical method for analysing such inhomogeneous phenomena we have robustly demonstrated that whilst drumlins can be randomly distributed (7% of drumlins analysed; 26% of flow-sets), their most common state (93% of drumlins analysed; 74% of flow-sets) is non-random. They exhibit a strong and statistically significant signal of regularity across lengths scales of 100 – 1200 m indicating that inter-drumlin interactions occur over this scale range. Regularity also occurs between 1200 to 10,000 m, but some of this might arise from regularity in the spacing of clusters in addition to longer range inter-drumlin interactions. In our database the finding of regularity holds across different ice sheets, countries, and across a variety of substrate geologies. Because of this, and the size of the database, we suggest regularity to be a near ubiquitous property of drumlins. Given that it took ourselves and many others so long to realise that the gaps in drumlin fields (due for example to lack of sediment and bedrock outcropping)

needed accounting for we now wish we had paid more attention to the words of Menzies in his review (1979, p338); 'It must be further noted that in most studies of drumlin spacing, little or no account is taken of other influencing structures such as roches moutonees or bedrock protrusions [i.e. gaps]. It is the writer's opinion that such influencing features are equally important in such morphometric studies.'

Although the spatial statistical analysis used in this paper demonstrates strong regularity the method says nothing about the spatial structure of the inherent organisation (i.e. how the pattern is organised). This is because the analysis works by assessing the departure from randomness using circles - increasingly long radii - from each drumlin. So we know that regularity exists but not the form it takes. Connecting this finding with the observations made earlier (albeit less quantitative or secure) about the transverse banding, we interpret that these are likely a large influence on the measured regularity, but also that lateral spacing contributes. The idealised form of spatial organisation of drumlins can thus be depicted as in Figure 13. We interpret that it is the higher density zones (drumlin patches) that misled some of the earlier statistical analyses of drumlin fields into concluding that clustering is the dominant signal. Well yes, clustering exists in drumlin fields but likely arises from antecedent variations in sediment thickness rather than from the drumlin-forming process. Once this is accounted for the drumlin-to-drumlin positioning is mostly regular.

The regularity of drumlin positioning demonstrates that they are a patterned phenomenon; a pattern being defined as a discernible regularity in which elements are repeated at specific length-scales. This finding is important as it should guide the quest for understanding the processes of drumlin formation and for evaluation of existing hypotheses. It says that a 'site-specific' landform view that considers each drumlin as a separate individual formed at a specific site and essentially independent of others, is not appropriate for most drumlin fields (but see later for rock-cored drumlins etc). Arguably it might be better to consider the drumlin array or field as the primary phenomenon and to regard them as bedforms *sensu stricto* (cf. Menzies and Rose, 1987), characterised simply by amplitude, relief and orientation. In this view the definition of drumlin boundaries, as in most studies including this are somewhat arbitrary and unnecessary. Ripples on a beach for example are not mapped and analysed individually but treated as a patch of patterned and bumpy bedforms with a length-scale, relief and orientation. Rather than speaking of drumlins perhaps it would be better to call the phenomenon drumlinised terrain.

When patterns are found in Nature they are usually organised by a process (Ball, 1999). Some of the most striking natural patterns are geomorphological (Werner, 1999) or biological (Koch and Meinhardt 1994). Exemplars include dunes (Kocurek and Ewing, 2005), freeze-thaw patterned ground (Kessler and Werner, 2003) river meander networks (Hallet, 1990), the fronds on a fern, and leaf positioning on a tree. It seems that all patterns in Nature must come into being by

interactions between elements as they evolve (grow or shrink). They are not 'printed' as a pattern but evolve into one. Elements of the pattern somehow 'know' of other elements to control their spacing, similar to geese in flight where the flock self-organises into an *en echelon* pattern by each individual keeping a set distance and angle from just one neighbour. We suggest that such self-organisation is intrinsic to the drumlinising process with interactions occurring between bumps at a kilometre scale; they must interact with each other whilst evolving. The spatial organisation of drumlins can therefore be regarded as an emergent property of the processes at work.

In Figure 14A we suggest that patterning of drumlins might arise from local self-organisation, analogous to how regularly-spaced cusps form on a beach (Werner and Fink, 1993). It is thought that a slight depression encourages flow acceleration of water and increased erosion to create embayments with deposition at the periphery and with regularity arising from such competition between adjacent embayments and the smoothing imparted at the scale of the beach. The point being that there is a positive feedback between sediment mobility and relief. The drumlin model of differential erosion by regelation-infiltration with variations mediated by effective pressure (Iverson, 2000) seems to be an example of this type.

In other pattern-forming geomorphological systems, such as ripples and dunes, progress has been made with explanations rooted in non-linear dynamics and

complex systems (Kocurek *et al.*, 2010; Murray *et al.*, 2014). Spontaneous development of a pattern from a system of mobile material with coupling between the fluid flow, sediment flux, and the bed shape, is usually taken to arise from instability in the system that amplifies tiny disturbances. Infinitesimally small perturbations at the bed grow at a preferred wavelength (spacing) and instability analysis predicts the wavelength at which bumps should grow, generating repeated culminations with regularity at this preferred scale (Figure 14C). One possibility then is that the scale of regularity we have identified in our database is a consequence of the wavelength of preferred growth in an unstable system. Examples of drumlin models of this type include the Hindmarsh-Fowler instability in the coupled flow of ice and sediment (Hindmarsh, 1998; Fowler, 2000), a thermo-mechanical instability in a partially frozen till substrate (Hooke and Medford, 2013) or as intrinsically required in the mega-flood hypothesis (Shaw, 2002). The latest version of the Hindmarsh-Fowler theory (Fowler and Chapwanya 2014) includes subglacial water flow within an upper layer of water-saturated sediments (i.e. more realistic than a thin water film across which stress cannot be transmitted) and this addition results in sediment erosion by water flow breaking up subglacial ribs to yield laterally-separated bumps resembling drumlins. Outputs from the numerical model estimate wavelengths consistent with measured drumlin dimensions (Clark *et al.* 2009) and the regularity that we report here. But there is some naivety in this view, in that it is not known if there is a mechanism that stabilises the bumps at this wavelength, rather than allowing them to develop further. There are reasons for expecting further evolution to

occur, and the prominent transverse banding found in drumlin fields might be an example of a coarsening process. From analysis of other geomorphological systems it is considered that finite-amplitude pattern formation could stabilise at the preferred wavelengths predicted by the instability analysis or, alternatively, as the bumps migrate they could collide and coalesce (Figure 14B) with such interactions (coarsening) resulting in regularity at an entirely different scale (cf. Kocurek *et al.*, 2010; Murray *et al.*, 2014).

Field measurements of recently formed drumlins in Iceland (McCracken *et al.* 2016) appear to be inconsistent with the Hindmarsh-Fowler instability theory (Fowler and Chapwanya 2014). If the identified instability mechanisms are not the underlying cause of drumlins it could be that individual drumlins initiate at randomly occurring locations, say triggered by stiff patches of sediment (although see Menzies *et al.*, 2016; Hart 1997), and then grow in dimensions over time such that they eventually meet and interact with their neighbours. Or they might migrate and interact by collisions (Figure 14B). Regularity would then somehow emerge from these meso-scale interactions.

Our demonstration that patterning exists across extensive areas known to have variations in sediment characteristics imply that such variations are of second-order importance to the process of drumlin formation. The concept of equifinality generates a problem of terminology, in that whilst drumlin-arrays (local patches of self-organised drumlins) arise from interactions as they evolve, it is also true that

shapes resembling drumlins (call them drumlins if you like) may form as 'site-specific landforms', say by sediment smeared around a bedrock protuberance. The latter do not falsify the finding of regularity, they just add some disorder (defects) to the pattern, and as proposed elsewhere (Clark, 2010) such elements (obstacle-drumlins and drumlin-clones) are likely to be different from the more ubiquitous emergent drumlins. We predict that emergent drumlins formed in thick drift (>15 m) exhibit stronger regularity than those in thin drift (<5 m) because in the latter case many of the forms will merely be obstacle-drumlins pinned to underlying bedrock protuberances, and which might be randomly distributed.

Ribbed moraine and MSGs have already been found to exhibit strongly preferred wavelengths (Dunlop and Clark, 2006; Spagnolo *et al.*, 2014) suggesting that they are also patterned phenomena requiring an explanation of their regular spacing. This patterning similarity between ribbed moraine, MSG and drumlins along with the finding that they vary in a continuum of shape and scale (Ely *et al.* 2016) adds credence to idea that they are genetically-related rather than separate manifestations formed by different processes.

In addition to the question of how drumlins form, the question of why they form has rarely been addressed (Smalley *et al.*, 2000). In considering why subglacial basal shear stresses are typically found to occupy a narrow physical range, Lliboutry (1958) hinted at an answer to both questions. He suggested that ice flow seeks to modify the bed roughness, smoothing out overly rough beds to

reduce friction, but when the bed is too smooth, drumlins are produced to provide resistance to flow. More work is required to explore interactions between roughness and flow. The ultimate goal is a combined understanding of how subglacial conditions yield a bumpy bed and the extent to which this roughness modulates ice flow velocity. That drumlins, along with MSGL and ribbed moraine, are found to occupy a set scale and pattern highlights the potential for using these predictable emergent scales of roughness to help inform the next-generation of ‘sliding laws’ to improve modelling of ice sheet and ice stream flow (Ritz et al. 2015).

Conclusions

Recognising that drumlin fields are patchy and contain gaps and choosing an appropriate statistical method for analysing such an inhomogeneous distribution we have shown that whilst drumlins are sometimes randomly placed, their main state is not random. They exhibit a strong and statistically significant signal of regularity across scales of 100 – 1200 m and possibly also up to 10,000 m in range. Because this finding holds across different ice sheets and a variety of substrate geologies we conclude that regularity is a ubiquitous tendency within drumlin fields. This patterning demonstrates spatial self-organisation in the bedforming process with drumlins as a specific emergent manifestation of subglacial sediment mobility. Elements of the pattern must have some physical interplay with other elements to control their spacing and so kilometre-scale interactions between drumlins must occur as they evolve, or interactions may

618 arise as a consequence of growth or migration. Hypotheses or models that can
619 explain the spacing of drumlins are required. We highlight three suggestions for
620 such self-organisation that are not mutually exclusive: instability in the coupling of
621 flow - sediment flux - bed shape; local feedback between sediment mobility and
622 relief; coarsening by growth or migration. Numerical modelling tracking bump
623 evolution beyond the initial instability phase (Chapwanya *et al.*, 2011) to explore
624 coarsening might prove to be fruitful, and further observations of bedforms
625 evolving beneath ice sheets (e.g. King *et al.*, 2009; 2016; Johnson *et al.*, 2010)
626 should lead to improved understanding.

627
628 That drumlins are now known to be patterned landforms, along with MSGL and
629 ribbed moraine, and that these have been found to vary in a continuum of shape
630 and scale (Ely *et al.* 2016) suggests that a unifying theory might be appropriate to
631 explain these subglacial bedforms. In such a view these named types are merely
632 emergent variants of the same phenomenon, whose differences arise from some
633 key parameter, such as ice velocity, cumulative basal sliding distance, sediment
634 thickness or from histories of interaction as they grow or migrate.

635 636 **Acknowledgements**

637 This work was supported by NERC grant NE/D011175/1 led by CDC. JCE was
638 generously supported by a PhD scholarship from Kathy and Chris Denison.
639 Andrew Fowler and Richard Hindmarsh are thanked for numerous illuminating
640 and otherwise conversations on the topic. David Unwin and Clive Anderson are

641 thanked for their pointers regarding spatial analysis and R software. We are
642 grateful to referees for helping improve the paper.

643

644 **References**

645 Aario, R. 1977. Classification and terminology of morainic landforms in Finland.

646 *Boreas*, **6(2)**, 87-100.

647 Armstrong JE, Tipper HW. 1948. Glaciation in north-central British Columbia.

648 *American Journal of Science* **246** : 283–310. DOI: 10.2475/ajs.246.5.283

649 Baddeley A, Diggle PJ, Hardegen A, Lawrence T, Milne RK, Nair G. 2014. On tests

650 of spatial pattern based on simulation envelopes. *Ecological monographs* **84** :

651 477–489.

652 Baddeley AJ, Møller J, Waagepetersen R. 2000. Non- and semi-parametric

653 estimation of interaction in inhomogeneous point patterns. *Statistica Neerlandica*

654 54 : 329–350. DOI: 10.1111/1467-9574.00144

655 Baddeley AJ., Rubak E., Turner R. (2015). Spatial Point Patterns, Methodology and

656 Application with R. CRC Press, Boca Raton, USA.

657 Ball P. 1999. The Self-Made Tapestry: Pattern formation in Nature. Oxford. Oxford

658 University Press.

659 Baranowski S. 1969. Some remarks on the origin of drumlins. *Geographia Polonica*,

660 **17**, 197-208.

661 Baranowski S. 1977. Regularity of drumlin distribution and the origin of their

662 formation. *Studia Geologica Polonica* **52** : 53–58.

663 Barchyn, T.E., Dowling, T.P.F., Stokes, C.R., Hugenholtz, C.H., 2016. Subglacial
664 bed form morphology controlled by ice speed and sediment thickness.
665 *Geophysical Research Letters*, **43**: 2016GL069558. doi:10.1002/2016GL069558

666 Boots BN, Burns RK. 1984. Analyzing the spatial distribution of drumlins: a two-
667 phase mosaic approach. *Journal of Glaciology* 30 : 302–307.

668 Brooke Benjamin T. 1957. Wave formation in laminar flow down an inclined plane.
669 *Journal of Fluid Mechanics* **2** : 554–573. DOI: 10.1017/S0022112057000373

670 Carl JD. 1978. Ribbed moraine–drumlin transition belt, St. Lawrence Valley, New
671 York. *Geology* **6** : 562–566.

672 Chapwanya M, Clark CD, Fowler AC. 2011. Numerical computations of a theoretical
673 model of ribbed moraine formation. *Earth Surface Processes and Landforms* **36** :
674 1105–1112. DOI: 10.1002/esp.2138

675 Clapperton CM. 1989. Asymmetrical drumlins in patagonia, chile. *Sedimentary*
676 *Geology* **62** : 387–398.

677 Clark CD. 1993. Mega-scale glacial lineations and cross-cutting ice-flow landforms.
678 *Earth Surface Processes and Landforms* **18** : 1–29. DOI:
679 10.1002/esp.3290180102

680 Clark CD. 1999. Glaciodynamic context of subglacial bedform generation and
681 preservation. *Annals of Glaciology* **28** : 23–32. DOI:
682 10.3189/172756499781821832

683 Clark CD. 2010. Emergent drumlins and their clones: from till dilatancy to flow
684 instabilities. *Journal of Glaciology* **56** : 1011–1025. DOI:
685 10.3189/002214311796406068

686 Clark CD, Wilson C. 1994. Spatial analysis of lineaments. *Computers &*
687 *Geosciences* **20** : 1237–1258. DOI: 10.1016/0098-3004(94)90073-6

688 Clark CD, Hughes ALC, Greenwood SL, Spagnolo M, Ng FSL. 2009. Size and
689 shape characteristics of drumlins, derived from a large sample, and associated
690 scaling laws. *Quaternary Science Reviews* **28** : 677–692. DOI:
691 10.1016/j.quascirev.2008.08.035

692 Couteron P, Seghier J, Chadœuf J. 2003. A test for spatial relationships between
693 neighbouring plants in plots of heterogeneous plant density. *Journal of vegetation*
694 *science: official organ of the International Association for Vegetation Science* **14** :
695 163–172. DOI: 10.1111/j.1654-1103.2003.tb02141.x

696 Dixon PM. 2012. Ripley's K function. In *Encyclopedia of Enviornmetrics*, , El-
697 Shaarawi A H And (ed). Chichester, John Wiley & Sons, Ltd.; 1796–1803.

698 Dunlop P, Clark CD, Hindmarsh RCA. 2008. Bed Ribbing Instability Explanation:
699 Testing a numerical model of ribbed moraine formation arising from coupled flow
700 of ice and subglacial sediment. *Journal of Geophysical Research* **113** : F03005.
701 DOI: 10.1029/2007JF000954

702 Dunlop P, Clark CD. 2006. The morphological characteristics of ribbed moraine.
703 *Quaternary Science Reviews* **25** : 1668–1691. DOI:
704 10.1016/j.quascirev.2006.01.002

705 Ely, J.C. 2015. Flow Signatures on the Bed and the Surface of Ice Sheets.
 706 Unpublished PhD thesis, University of Sheffield, pp 300.

707 Ely JC, Clark CD, Spagnolo M, Stokes CR, Greenwood SL, Hughes ALC, Dunlop P,
 708 Hess D. 2016. Do subglacial bedforms comprise a size and shape continuum?
 709 *Geomorphology* **257** : 108–119. DOI: 10.1016/j.geomorph.2016.01.001

710 Eyles N, Putkinen N, Sookhan S, Arbelaez-Moreno L. 2016. Erosional origin of
 711 drumlins and megaridges. *Sedimentary Geology* DOI:
 712 10.1016/j.sedgeo.2016.01.006

713 Fowler AC, Chapwanya M. 2014. An instability theory for the formation of ribbed
 714 moraine, drumlins and mega-scale glacial lineations. *Proceedings of the Royal*
 715 *Society A* **470** : 20140185. DOI: 10.1098/rspa.2014.0185

716 Fowler AC, Spagnolo M, Clark CD, Stokes CR, Hughes ALC, Dunlop P. 2013. On
 717 the size and shape of drumlins. *GEM - International Journal on Geomathematics*
 718 **4** : 155–165. DOI: 10.1007/s13137-013-0050-0

719 Fowler AC. 2000. An instability mechanism for drumlin formation. Geological
 720 Society, London, Special Publications **176** : 307–319. DOI:
 721 10.1144/GSL.SP.2000.176.01.23

722 Francek MA. 1991. A spatial perspective on the New York drumlin field. *Physical*
 723 *Geography* **12** : 1–18. DOI: 10.1080/02723646.1991.10642415

724 Gao X, Narteau C, Rozier O, du Pont SC. 2015. Phase diagrams of dune shape
 725 and orientation depending on sand availability. *Scientific Reports* **5** : 14677. DOI:
 726 10.1038/srep14677

727 Greenwood SL, Clark CD. 2010. The sensitivity of subglacial bedform size and
 728 distribution to substrate lithological control. *Sedimentary Geology* **232** : 130–144.
 729 DOI: 10.1016/j.sedgeo.2010.01.009

730 Greenwood SL, Kleman J. 2010. Glacial landforms of extreme size in the Keewatin
 731 sector of the Laurentide Ice Sheet. *Quaternary science reviews* **29** : 1894–1910.
 732 DOI: 10.1016/j.quascirev.2010.04.010

733 Hall J. 1815. On the revolutions of the Earth's surface. *Transactions of the Royal*
 734 *Society of Edinburgh* **7** : 169–184.

735 Hallet B. 1990. Spatial self-organization in geomorphology: from periodic bedforms
 736 and patterned ground to scale-invariant topography. *Earth-Science Reviews* **29** :
 737 57–75. DOI: 10.1016/0012-8252(90)90028-t

738 Hart JK. 1997. The relationship between drumlins and other forms of subglacial
 739 glaciotectionic deformation. *Quaternary Science Reviews* **16** : 93–107. DOI:
 740 10.1016/s0277-3791(96)00023-6

741 Hättestrand C. 1997 Ribbed moraines in Sweden – distribution, pattern and
 742 palaeoglaciological implications. *Sedimentary Geology* **111**: 41-56.

743 Hill AR. 1973. The distribution of drumlins in County Down, Ireland. *Annals of the*
 744 *Association of American Geographers*. Association of American Geographers **63**
 745 : 226–240.

746 Hindmarsh, R. C. A. 1998. Drumlinization and drumlin-forming instabilities: viscous
 747 till mechanisms. *Journal of Glaciology*, **44**(147), 293-314.

748 Hooke RL, Medford A. 2013. Are drumlins a product of a thermo-mechanical
 749 instability? *Quaternary Research* **79** : 458–464. DOI:
 750 10.1016/j.yqres.2012.12.002

751 Hughes ALC, Clark CD, Jordan CJ. 2010. Subglacial bedforms of the last British Ice
 752 Sheet. *Journal of Maps* **6** : 543–563. DOI: 10.4113/jom.2010.1111

753 Hughes ALC, Clark CD, Jordan CJ. 2014. Flow-pattern evolution of the last British
 754 Ice Sheet. *Quaternary Science Reviews* **89** ; 148-168. DOI:
 755 10.1016/j.quascirev.2014.02.002

756 Illian J., Penttinen A., Stoyan H., Stoyan, D. 2008. Statistical Analysis and Modelling
 757 of Spatial Point Patterns. John Wiley & Sons Ltd., Chichester, UK.

758 Iverson NR. 2000. Sediment entrainment by a soft-bedded glacier: a model based
 759 on regelation into the bed. *Earth Surface Processes and Landforms* **25** : 881–
 760 893. DOI: 10.1002/1096-9837(200008)25:8<881::AID-ESP105>3.0.CO;2-Y

761 Jauhiainen F. 1975. Morphometric analysis of drumlin fields in northern Central
 762 Europe. *Boreas* **4** : 219–230. DOI: 10.1111/j.1502-3885.1975.tb00691.x

763 Johnson MD, Schomacker A, Benediktsson ÍÖ, Geiger AJ, Ferguson A, Ingólfsson
 764 Ó. 2010. Active drumlin field revealed at the margin of Múlajökull, Iceland: A
 765 surge-type glacier. *Geology* **38** : 943–946. DOI: 10.1130/G31371.1

766 Jónsson SA, Schomacker A, Benediktsson ÍÖ, Ingólfsson Ó, Johnson MD. 2014.
 767 The drumlin field and the geomorphology of the Múlajökull surge-type glacier,
 768 central Iceland. *Geomorphology* **207** : 213–220. DOI:
 769 10.1016/j.geomorph.2013.11.007

770 Kessler MA, Werner BT. 2003. Self-organization of sorted patterned ground.
 771 *Science* **299** : 380–383. DOI: 10.1126/science.1077309

772 King EC, Hindmarsh RCA, Stokes CR. 2009. Formation of mega-scale glacial
 773 lineations observed beneath a West Antarctic ice stream. *Nature Geoscience* **2** :
 774 585–588. DOI: 10.1038/ngeo581

775 King, E.C., Pritchard, H.D., Smith, A.M., 2016. Subglacial landforms beneath
 776 Rutford Ice Stream, Antarctica: detailed bed topography from ice-penetrating
 777 radar. *Earth System Science Data* **8**: 151- 158.

778 Koch AJ, Meinhardt H. 1994. Biological pattern formation: from basic mechanisms
 779 to complex structures. *Reviews of Modern Physics* **66** : 1481–1507. DOI:
 780 10.1103/RevModPhys.66.1481

781 Kocurek G, Ewing RC. 2005. Aeolian dune field self-organization--implications for
 782 the formation of simple versus complex dune-field patterns. *Geomorphology* **72** :
 783 94–105. [

784 Kocurek G, Ewing RC, Mohrig D. 2010. How do bedform patterns arise? New views
 785 on the role of bedform interactions within a set of boundary conditions. *Earth*
 786 *Surface Processes and Landforms* **35** : 51–63. DOI: 10.1002/esp.1913

787 Kulldorff M. 2006. Tests of Spatial Randomness Adjusted for an Inhomogeneity: A
 788 General Framework. *Journal of the American Statistical Association* **101** : 1289–
 789 1305.

790 Lliboutry L. 1958. Contribution à la théorie du frottement du glacier sur son lit. C. R.
 791 *Hebd. Séances Acad. Sci.* **247** : 318–320.

792 McCracken, R.G., Iverson, N.R., Benediktsson, Í.Ö., Schomacker, A., Zoet, L.K.,
 793 Johnson, M.D., Hooyer, T.S., Ingólfsson, Ó., 2016. Origin of the active drumlin
 794 field at Múlajökull, Iceland: New insights from till shear and consolidation
 795 patterns. *Quaternary Science Reviews* **148**: 243–260.

796 MacLachlan JC, Eyles CH. 2013. Quantitative geomorphological analysis of drumlins
 797 in the Peterborough drumlin field, Ontario, Canada. *Geografiska Annaler. Series*
 798 *A, Physical Geography* **95** : 125–144.

799 Menzies J. 1979. A review of the literature on the formation and location of
 800 drumlins. *Earth-Science Reviews* **14** : 315–359.

801 Menzies J, Rose J. 1987. Drumlins-trends and perspectives. *Episodes* **10** : 29–31.

802 Menzies J, Hess DP, Rice JM, Wagner KG, Ravier E. 2016. A case study in the
 803 New York Drumlin Field, an investigation using microsedimentology, resulting in
 804 the refinement of a theory of drumlin formation. *Sedimentary Geology* **338** : 84–
 805 96. DOI: 10.1016/j.sedgeo.2016.01.017

806 Murray AB, Goldstein EB, Coco G. 2014. The shape of patterns to come: from initial
 807 formation to long-term evolution. *Earth Surface Processes and Landforms* **39** :
 808 62–70.

809 Myllymäki M, Mrkvička T, Grabarnik P, Seijo H, Hahn U. 2016. Global envelope
 810 tests for spatial processes. *Journal of the Royal Statistical Society, Series B*.
 811 DOI: 10.1111/rssb.12172

812 O'Sullivan D, Unwin DJ. 2010. *Geographic Information Analysis and Spatial Data*.
813 Geographic Information Analysis, Second Edition, John Wiley & Sons, Inc. ,
814 Hoboken, NJ, USA.

815 Patterson CJ, Hooke, RLeb. 1995. Physical environment of drumlin formation.
816 *Journal of Glaciology* **41** : 30–38.

817 Perry GLW, Miller BP, Enright NJ. 2006. A comparison of methods for the statistical
818 analysis of spatial point patterns in plant ecology. *Plant Ecology* **187** : 59–82.
819 DOI: 10.1007/s11258-006-9133-4

820 Reed B, Galvin CJ, Miller JP. 1962. Some aspects of drumlin geometry. *American*
821 *journal of Science* **260** : 200–210. DOI: 10.2475/ajs.260.3.200

822 Ripley B. 1988. *Statistical Inference for Spatial Processes*. Cambridge University
823 Press.

824 Ripley, B D and Rasson, J P. 1977. Finding the edge of a Poisson forest. *Journal of*
825 *Applied Probability* **14** : 483 – 491.

826 Ritz C, Edwards TL, Durand G, Payne AJ, Peyaud V, Hindmarsh RCA. 2015.
827 Potential sea-level rise from Antarctic ice-sheet instability constrained by
828 observations. *Nature* **528** : 115–118. DOI: 10.1038/nature16147

829 Rose J, Letzer JM. 1977. Superimposed drumlins. *Journal of Glaciology* **18** : 471–
830 480.

831 Rose, J. 1987. Drumlins as part of a glacier bedform continuum. In Menzies, J. and
832 Rose, J. (eds), Drumlin Symposium. Rotterdam, Balkema, 103-116.

833 Shaw J, Kvill D. 1984. A glaciofluvial origin for drumlins of the Livingstone Lake
834 area, Saskatchewan. *Canadian Journal of Earth Sciences* **21** : 1442–1459. DOI:
835 10.1139/e84-150

836 Shaw J. 2002. The meltwater hypothesis for subglacial bedforms. *Quaternary*
837 *International*: **90**: 5–22. DOI: 10.1016/S1040-6182(01)00089-1

838 Smalley I, Warburton J. 1994. The shape of drumlins, their distribution in drumlin
839 fields, and the nature of the sub-ice shaping forces. *Sedimentary Geology* **91** :
840 241–252. DOI: 10.1016/0037-0738(94)90132-5

841 Smalley IJ, Lu P, Jefferson IF. 2000. The golf ball model and the purpose of drumlin
842 formation. *Studia Quaternaria* **17** : 29–33.

843 Smalley IJ, Unwin DJ. 1968. The formation and shape of drumlins and their
844 distribution and orientation in drumlin fields. *Journal of Glaciology* **7** : 377–390.

845 Spagnolo M, Clark CD, Hughes ALC. 2012. Drumlin relief. *Geomorphology* **153–**
846 **154** : 179–191. DOI: 10.1016/j.geomorph.2012.02.023

847 Spagnolo M, Clark CD, Ely JC, Stokes CR, Anderson JB, Andreassen K, Graham
848 AGC, King EC. 2014. Size, shape and spatial arrangement of mega-scale glacial
849 lineations from a large and diverse dataset. *Earth Surface Processes and*
850 *Landforms* **39** : 1432–1448. DOI: 10.1002/esp.3532

851 Stokes CR, Spagnolo M, Clark CD. 2011. The composition and internal structure of
852 drumlins: complexity, commonality, and implications for a unifying theory of their
853 formation. *Earth-Science Reviews* **107**, 398-422.

854 Trenhaile AS. 1971. Drumlins: Their distribution, orientation and morphology.
 855 *Canadian Geographer / Le Géographe canadien* **15** : 113–126. DOI:
 856 10.1111/j.1541-0064.1971.tb00147.x

857 Unwin D. 1997. *GIS and spatial statistical analysis*. Geographic Information
 858 Research: Bridging the Atlantic (M. Craglia and H. Couclelis, eds.). London:
 859 Taylor & Francis : 399–411.

860 Vernon, P. 1966. Drumlins and Pleistocene ice flow over the Ards Peninsula,
 861 Strangford Lough area, County Down, Ireland. *Journal of Glaciology*, **6** (45), 401-
 862 409.

863 Werner BT, Fink TM. 1993. Beach cusps as self-organized patterns. *Science* **260** :
 864 968–971. DOI: 10.1126/science.260.5110.968

865 Werner BT. 1999. Complexity in natural landform patterns. *Science* **284** : 102–104.


866 Werth E. (1909) Eine Drumlinlandschaft und Rirmenseen SE von Posen,
 867 *Zitschr.d.d. Geol. Ges.*,61.

868

Citation	Number of drumlins investigated	Location	Statistical method applied	Concluded type of organisation
Reed et al., 1962	204 across two fields	USA	Spacing measurements	Preferred spacing
Smalley and Unwin, 1968	203 across four fields	Ireland	Clark and Evans (1954) nearest neighbour statistic	Random
Vernon, 1966	733	Ireland and England	Nearest neighbour distances (NNd)	Variable spacing
Baranowski, 1969	1054 across two fields	USA and Poland	Nearest neighbour measurements and transvers profiles	Regular spacing
Trenhaile, 1971	~ 7000 across several fields	Canada	Comparison to two random distribution models, and order-rank nearest neighbour analysis	Between random and uniformly spaced
Hill, 1973	~3900	Ireland	Nearest neighbour measurements, trend surface analysis and cumulative frequency distributions of NNd	Non-randomly distributed, mostly clustered. Preferred spacing in NNd
Jauhiainen, 1975	1137 across twelve fields	Northern Europe	Clark and Evans (1954) nearest neighbour statistic	Clustered to random
Boots and Burns, 1984	102 across two fields	Canada and England	Two phase mosaic	Random
Francek, 1991	4054	USA	Clark and Evans (1954) nearest neighbour statistic	Mostly random, some tendency toward clustering
MacLachlan and Eyles, 2013	~3000	Canada	Ripley's K-function	Clustered

870


871 **Table 1. Summary of investigations on the type of spatial organisation of**
872 **drumlins and which highlights the variety of findings from random,**
873 **clustered to regular.**



875

876 Figure 1. Types of drumlin spatial organisation. A) En echelon arrangement with
 877 the next lateral drumlin positioned slightly downstream (like geese in flight, with
 878 each bird positioned to avoid turbulence from an upwind flock member). B)
 879 Transverse banding in drumlin densities. C) Packed versus widely-spaced
 880 distribution. D) Random, clustered or regularly spaced positioning within a flow-
 881 set, the latter having a preferred spacing expressed as a single narrow peak in
 882 the frequency distribution. In spatial statistics an homogeneous distribution of
 883 features is taken to be that they are uniformly distributed across an area (i.e. no
 884 large gaps). As argued in the text and apparent here and in Fig 2. we show that
 885 drumlins are more appropriately considered as being inhomogeneously
 886 distributed in that they often occur in patches with large gaps between them.

887



888

889 Figure 2. Drumlins appearing as elongated bumps depicted here by digital
 890 elevation models. a) Drumlins in Ireland (Co. Leitrim) with a widely-spaced
 891 arrangement and with seemingly regular spacing and some en echelon
 892 positioning. Black line marks profile in (c). b) More tightly packed drumlins in
 893 England (Yorkshire Dales) and with elements of transverse banding evident in
 894 drumlin positioning, and with gaps in the drumlin field corresponding to higher
 895 elevations with thin or absent sediment cover. c) Elevation profile across drumlins
 896 of ca. 20 m relief, from profile (black line) in a.



897

898 Figure 3. Location of drumlins used in the analyses. 100 flow-sets (representing
 899 different flow events) of the last British Ice Sheet, from Hughes et al. (2010;
 900 2014) and comprising 36,222 drumlins, plus additional flow-sets from Alta,
 901 Norway (n = 1,483) and Ungava Bay, Canada (n = 5,903), making a total of
 902 43,608. Flow-sets with a low number of drumlins (<30) were excluded from the
 903 analyses leaving a sample of 42,488 drumlins.





Figure 4. Comparison of transverse banding evident in a real drumlin field (a; Leitrim Ireland) with a simulated one (b) using ellipses of the same size and orientation but positioned randomly. A few red lines are drawn in to illustrate banding, many more are visible. Note that in b) some transverse banding and echelon positioning is evident - purely by chance - but is less prevalent and it is much rarer to find similarly-sized drumlins comprising a band.

912
913



914
915
916
917
918
919
920
921
922
923

Figure 5. Drumlin patchiness and Ripley's L analysis. a) A complete flow-set of drumlins (black points) and a small visibly homogeneous patch of 4000 x 5000 m (blue rectangle) b) Output from the L-function for the complete data set (black) and the patch (blue). Red dashed line is the expectation under complete spatial randomness. The whole flow-set shows some small scale regularity (dispersion), probably determined by drumlin size that quickly crosses into clustering but the small patch looks regular over a wider scale range, up to around 300 m.




Figure 6. The problem of patchiness (density variations) in drumlin fields for analysis of spatial organisation. The preference for drumlins to occur in patches of higher density amongst more sparsely distributed forms yields a clustering in the spatial arrangement, and with some gaps often arising from lack of sedimentary cover. a) Typical example from northern England (flowset 66 of Hughes et al., 2014) with large gaps and patches of varying density. In b) to illustrate a point we show several patches of graph paper (each with obvious regularity). Ripley's L function applied to both the drumlins (a) and the graph paper (b) yield clustering as the signal (c), and yet for the graph paper (b) we know that regularity actually exists. The inhomogeneous pairwise correlation function however, takes account of the inherent clustering and shows (d) that for both the drumlins and the graph paper there is a strong preference (attraction) for points to be spaced a certain distance apart. Clustering has not masked the regularity that exists.




Figure 7. Output from inhomogeneous pair-correlation function on flow-set 69, near Kirkby Lonsdale, Cumbria, England. Note the peak in attraction at approximately $r = 450$ m and the initial repulsion of points before this scale. Smoothing of the observed function in R is automatically performed. In order to highlight regions of the graph, tests at limited spatial scales (e.g. up to 4000 m) were also performed, and smoothing of functions altered accordingly. Here, the envelope is produced from 499 simulations of inhomogeneous CSR, yielding a p value of 0.002. The red line indicates the assumed value under CSR. The grey envelope is derived from the Monte-Carlo simulations of CSR and the black line is the function of the observed points.




Figure 8. Sensitivity analysis of the function $g_{inhom}(r)$ to different intensity search radii for flow-set 29. Envelopes of complete spatial randomness (grey) are significant to $p = 0.002$.




Figure 9. Of all British drumlins in our database 48% were found to belong to transverse bands suggesting that this style of spatial organisation is a common feature of drumlin flow-sets. Drumlins classified by our method as belonging to a band are depicted as black.





Figure 10. The spacing of drumlins. Frequency distribution of nearest neighbour distances (avoiding double-counting) between drumlin summits in the UK database ($n=36,222$). The uni-modal peak centred on 325 m is interpreted as indicating a clearly defined preference for the spacing of drumlins, suggestive - but not proving - the existence of patterned spatial organisation.

971



972
973
974
975
976
977
978
979
980
981
982
983
984

Figure 11. The inhomogeneous pairwise correlation function (PCF) for flow-set fs29 in the Southern Uplands of Scotland comprising 1473 drumlins. The function $g_{\text{inhom}}(r)$ is plotted against r , with increasing radii of analysis (in m) away from each drumlin. The black line plots the function for the drumlins which should be compared against the red line and grey envelope representing complete spatial randomness (CSR). The peak at 330 m and extending between 160 – 1250 m indicates a statistically significant demonstration of regularity (i.e. above grey envelope) in drumlin positioning across this length scale; ‘attraction’ of points to this distance has occurred, typified by a regular spatial arrangement. Values plotting within the grey envelope are not significantly different from a random spatial arrangement.




Figure 12. Demonstration of regular patterning in drumlin arrangement. All 72 flow-sets analysed (with $n > 30$) displayed on a log-log plot illustrating that the main signal is of regular patterning. The majority of flow-sets (blue lines; 74% of total) are found to have a statistically significant demonstration of regularity, representing 93% of the drumlin sample ($n = 42,488$). The green lines are for those flow-sets that did not exhibit regularity (26% of the sample), approximating either complete spatial randomness or clustering. The black line is for flow-set 29, from Fig 11. In the middle panel, black dots record the position of the first prominent peak of each curve (main signal of regularity at this scale) and blue dots mark where the signals decline into CSR. These provide the scale ranges for our conclusions in the panel beneath.




Figure 13. Drumlins are not randomly distributed; rather they are a patterned phenomenon that we idealise in this cartoon. They mostly occur in patches of higher density with internal regularity and banding amongst more sparsely distributed drumlins and with numerous gaps evident in the overall drumlin field. The gaps might be due to thin sediment cover (or bedrock exposed in places) or inappropriate sediment properties.




Figure 14. Suggestions of self-organisation that might individually, or in some combination, explain patterning of drumlinised terrain. In A, adjacent bumps interact and from this emerges a field-scale pattern with characteristic spacing, whereas as in B, a perhaps random distribution of bumps self-organise into a pattern by collisions arising from migration and growth. In C, instability occurs with the spacing controlled by the wavelength of the maximum growth rate. In this case simple stabilisation occurs at this wavelength, but it could be that migration and coarsening also occurs (i.e. as per B) after the initial perturbation.